

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07D 305/14	A1	(11) International Publication Number: WO 93/24476
		(43) International Publication Date: 9 December 1993 (09.12.93)

(21) International Application Number: PCT/US93/05344

(22) International Filing Date: 4 June 1993 (04.06.93)

(30) Priority data:
07/893,500 4 June 1992 (04.06.92) US(60) Parent Application or Grant
(63) Related by Continuation
US
Filed on 07/893,500 (CON)
4 June 1992 (04.06.92)(71) Applicant (for all designated States except US): CLOVER
CONSOLIDATED, LIMITED [CH/CH]; 37, avenue de
Rumini, CH-1002 Lausanne (CH).


(72) Inventors; and

(75) Inventors/Applicants (for US only): DESAI, Neil, P. [IN/
US]; 847 Alandale Avenue, Los Angeles, CA 90036
(US). SOON-SHIONG, Patrick [US/US]; 12307 Doroth
y Street, Los Angeles, CA 90049 (US). SANDFORD,
Paul, A. [US/US]; 2822 Overland Avenue, Los Angeles,
CA 90064 (US).(74) Agent: REITER, Stephen, E.; Pretty, Schroeder, Bruegge-
mann & Clark, 444 South Flower Street, Suite 2000, Los
Angeles, CA 90071 (US).(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ,
DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU,
MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE,
SK, UA, US, European patent (AT, BE, CH, DE, DK,
ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI
patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR,
NE, SN, TD, TG).

Published

With international search report.

(54) Title: WATER-SOLUBLE POLYMERIC CARRIERS FOR DRUG DELIVERY



(57) Abstract

In accordance with the present invention, there are provided polymeric drug delivery systems in which the drug is bound to a water-soluble polymer to provide a form of soluble drug delivery especially for those cases in which the drug by itself is water-insoluble. In particular, the drug taxol is covalently bound to water-soluble polyethylene glycols such as linear polyethylene glycols, branched polyethylene glycols, star polyethylene glycols, and branched copolymers of polyethylene glycols with other functional monomers to comprise a form of polymeric drug delivery. Also, cross-linked insoluble gels of these materials are prepared to serve as a form of implantable drug delivery.

ATTORNEY DOCKET NUMBER: 10177-191-999

SERIAL NUMBER: 10/603,115

REFERENCE: B120


BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07D 305/14		A1	(11) International Publication Number: WO 93/24476 (43) International Publication Date: 9 December 1993 (09.12.93)
(21) International Application Number: PCT/US93/05344 (22) International Filing Date: 4 June 1993 (04.06.93) (30) Priority data: 07/893,500 4 June 1992 (04.06.92) US (60) Parent Application or Grant (63) Related by Continuation US Filed on 07/893,500 (CON) 4 June 1992 (04.06.92) (71) Applicant (for all designated States except US): CLOVER CONSOLIDATED, LIMITED [CH/CH]; 37, avenue de Rumini, CH-1002 Lausanne (CH).		(72) Inventors; and (75) Inventors/Applicants (for US only) : DESAI, Neil, P. [IN/US]; 847 Alandale Avenue, Los Angeles, CA 90036 (US). SOON-SHIONG, Patrick [US/US]; 12307 Dorothy Street, Los Angeles, CA 90049 (US). SANDFORD, Paul, A. [US/US]; 2822 Overland Avenue, Los Angeles, CA 90064 (US). (74) Agent: REITER, Stephen, E.; Pretty, Schroeder, Brueggemann & Clark, 444 South Flower Street, Suite 2000, Los Angeles, CA 90071 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: **WATER-SOLUBLE POLYMERIC CARRIERS FOR DRUG DELIVERY**

(57) Abstract

In accordance with the present invention, there are provided polymeric drug delivery systems in which the drug is bound to a water-soluble polymer to provide a form of soluble drug delivery especially for those cases in which the drug by itself is water-insoluble. In particular, the drug taxol is covalently bound to water-soluble polyethylene glycols such as linear polyethylene glycols, branched polyethylene glycols, star polyethylene glycols, and branched copolymers of polyethylene glycols with other functional monomers to comprise a form of polymeric drug delivery. Also, cross-linked insoluble gels of these materials are prepared to serve as a form of implantable drug delivery.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

WATER-SOLUBLE POLYMERIC CARRIERS
FOR DRUG DELIVERY

FIELD OF THE INVENTION

The present invention relates to the drug delivery of taxol wherein the drug is chemically bound to a water-soluble polymeric or macromolecular carrier that renders the drug water-soluble. In another aspect, the present invention relates to water-soluble prodrugs of taxol that recover their biological activity when hydrolyzed from the carrier molecule. In a further aspect, the present invention relates to sustained drug delivery of taxol by hydrolysis from an implanted gel comprising the drug-carrier conjugate.

BACKGROUND OF THE INVENTION

Taxol is a natural product first isolated from the Pacific Yew tree, *Taxus brevifolia*, by Wani et al. (1971, *J. Am. Chem. Soc.* 93:2325). Among the antimitotic agents, taxol, which contains a diterpene carbon skeleton, exhibits a unique mode of action on microtubule proteins responsible for the formation of the mitotic spindle. In contrast with other antimitotic agents such as vinblastine or colchicine, which prevent the assembly of tubulin, taxol is the only plant product known to inhibit the depolymerization process of tubulin. This prevents the cell replication process and taxol has been shown to have significant antineoplastic and anticancer effects in drug-refractory ovarian cancer. Taxol has shown excellent antitumor activity in a wide variety of tumor models such as the B16 melanoma, L1210 leukemias, MX-1 mammary tumors, and CX-1 colon tumor xenografts. Several recent press releases have termed taxol as the new anticancer wonder-drug. The poor aqueous solubility of taxol has, however, remained a setback in human clinical trials, and currently used formulations require a cremaphore to solubilize the drug. The human clinical dose range is 200-500 mg and

requires about one liter of fluid given intravenously using the cremaphore. In phase I clinical trials taxol itself did not show excessive toxic effects but severe allergic reactions were caused by the emulsifiers administered to
5 solubilize the drug.

The general chemical structure of taxol is shown in Figure 1 in which $R_1 = R_2 = OH$. Potential sites for modification of the drug are at the hydroxyls on the 1, 7, and 2' carbon atoms. The 1-hydroxyl is sterically hindered
10 and nonreactive, the 2'-hydroxyl is the most reactive, followed by the 7-hydroxyl which is also sterically hindered. Thus the modification of taxol to increase its water-solubility has revolved around the modification of the 2'- and the 7-hydroxyls. Studies have reported that
15 the C-13 ester side chain and the 2'-hydroxyl group on the side chain are essential for biological activity. Mellado et al. (1984; *Biochem. Biophys. Res. Commun.* 124:329-336) have reported the synthesis of 2'-acetyl, 7-acetyl, and 2',7-diacetyl taxol. An acetyl at the 2' position resulted
20 in a loss in ability to promote microtubule assembly. Taxol and 7-acetyl taxol were similar in their ability to alter cell proliferation and microtubule polymerization. These observations suggest that the 2'- and 7-positions are suitable for structural modifications, the 2'-position as
25 a site for reversible derivatization (or formation of a prodrug) and the 7-position for analogue/prodrug modifications.

A number of chemically modified taxols with enhanced water-solubilities have been developed. Among
30 them are the sulfonated derivatives (Kingston et al., 1991, US Patent 5,059,699), and amino acid esters (Mathew et al., 1992; *J. Med. Chem.* 35:145-151) Which show significant biological activity. However, the delivery of taxol attached to a macromolecular or polymeric water-soluble
35 carrier has not been considered. Nathan et al. (1990;

Polymer Preprints 31:213-214) have described a polyethylene glycol (PEG) chain-extended with amino acids such as lysine, to produce a polymer which has pendant carboxylic acid groups that may be used to attach biologically active molecules. However, no mention is made of the immobilization of taxol, or the attachment of a water-insoluble drug to such a carrier in order to deliver it in a soluble form.

In the present invention, to deliver taxol in a water-soluble form we have used a water-soluble polymer to which the drug is bound, the resultant polymer-drug conjugate being soluble. Water-soluble polymers such as PEG, have been investigated extensively in recent years for use as nontoxic, biocompatible, protein repulsive, noninflammatory, and nonimmunogenic modifiers for drugs, proteins, enzymes, and surfaces of implanted materials. These characteristics have been variously attributed to a combination of properties of these polymers, e.g., nonionic character, water solubility, backbone flexibility, and volume exclusion effect in solution or when immobilized at a surface. The solubility of PEG in water as well as a number of common organic solvents facilitates its modification by a variety of chemical reactions and makes it amenable for binding water-insoluble or poorly water-soluble molecules and rendering them water-soluble.

The preparation of a reversible PEG-taxol derivative at the 2'- and/or 7-position on taxol serves as useful aqueous-soluble prodrug. A nonreversible PEG derivative on the 7-position of taxol serves as a useful water-soluble drug analogue.

Advantages of delivering the drug attached to a water-soluble polymer as described in the present invention are many fold. The number of drug molecules per polymer molecule can be controlled; the circulation time of the

drug can be varied by adjusting a number of variables including molecular weight of the polymeric carrier, the type of linkage between the drug and polymer, i.e., some linkages are hydrolyzed at much faster rates than others; large increases or decreases in blood levels of the drug may be avoided in favor of more gradual and sustained levels obtained by continuous release of the drug from a polymeric carrier; and the hydrolysis of the drug-polymer conjugate results in the formation of the original biologically active drug and the innocuous water-soluble polymer that is excreted from the body.

BRIEF SUMMARY OF THE INVENTION

The present invention relates to a method of drug delivery that utilizes water-soluble polymers as carriers for a drug. The delivery of drugs that are inherently insoluble or poorly soluble in an aqueous medium can be seriously impaired if the only suitable mode of delivery is by intravenous injection. The attachment of such drugs to water-soluble macromolecules that act as carriers can greatly benefit this problem and allow for intravenous, subcutaneous, or intramuscular delivery. Examples of poorly aqueous drugs that may benefit from this form of drug delivery are taxol, amphotericin B, etc. Examples of water-soluble polymers that may be used as carriers in such a system are polyethylene glycols (PEG), polyvinyl alcohol, polyhydroxyethyl methacrylate, polyacrylamide, polyacrylic acid, polyethyloxazoline, polyvinyl pyrrolidinone, and polysaccharides such as chitosan, alginates, hyaluronic acid, dextrans, etc.

In a preferred embodiment, the drug to be delivered is taxol, a naturally occurring diterpenoid which has been described as a potent antineoplastic and anticancer agent, and the polymeric water-soluble carrier is polyethylene glycol and derivatives thereof.

In another preferred embodiment, taxol is covalently linked to a PEG carboxylic acid derivative by an esterification at the 2'-position on the taxol side chain.

Another preferred embodiment involves the
5 esterification of taxol at the 2' position with succinic anhydride or glutaric anhydride followed by esterification with PEG to obtain the PEG-taxol derivative.

In yet another preferred embodiment, a multi-arm
10 'star' or 'branched' PEG is used as the carrier to increase the loading (number of drug molecules per carrier molecule) of taxol on the PEG.

In another preferred embodiment, an acrylate
15 derivative of PEG is copolymerized with acrylic acid to obtain a copolymer with a multiplicity of carboxyl functionalities that are sites for the attachment for taxol.

Another embodiment of the present invention is to
covalently attach taxol to a PEG-amine derivative by first
reacting taxol with carbonyldiimidazole followed by
20 reaction with PEG-amine to obtain a urethane linkage. This link is not readily hydrolyzable and such a derivative at the 2'-position interferes with the biological activity of taxol. It is therefore an embodiment of the present invention to produce such a PEG derivative at the 7-
25 position of taxol which retains its biological activity.

In another embodiment, the drug may be linked to
a star or branched PEG in which a part of the endgroups of
the PEG have been covalently linked to the drug while the
remainder are covalently linked to an unsaturated group
30 such as the acrylate group that may be polymerized in a free radical process to obtain a crosslinked polymer. The resultant crosslinked polymer, absorbs water in aqueous

medium and results in the formation of a hydrogel containing bound drug. This hydrogel may be implanted in a suitable location subcutaneously or intraperitoneally for sustained release of the drug by hydrolysis from the insoluble crosslinked carrier.

Thus it is a primary object of this invention to produce a derivative of taxol on a water-soluble macromolecule or polymer as a carrier that can be used for delivery of taxol in a soluble form.

It is a further object of the present invention to use a hydrolyzable linking group such as an ester to allow for the hydrolysis of the drug-polymer conjugate subsequent to delivery of the drug to form the original active drug and polymeric carrier.

It is yet another object of the invention that the drug produced upon hydrolysis retain its original biological activity and also the nonhydrolyzable derivative of the drug maintain its biological activity.

It is still a further object of the present invention to simplify the purification of the water-soluble conjugate by utilizing a polymeric carrier such as PEG that can be isolated by a simple precipitation.

DESCRIPTION OF THE DRAWINGS

Figure 1 is a general chemical structure for derivatives of Taxol where R_1 and R_2 are the sites of derivatization. The structure represents the unmodified Taxol molecule when $R_1 = R_2 = OH$.

DETAILED DESCRIPTION OF THE INVENTION

Water-soluble polymers such as PEG (Aldrich), and monomethoxy PEG (MPEG, Nippon Oil and Fats) were utilized to bind poorly aqueous-soluble drugs. Taxol (Sigma chemical) was the drug utilized for covalent linking to the carrier polymers. The 8-arm 'star' PEG polymer (MW 22800) was obtained from Macrochem Labs and acrylic acid from Aldrich. It should be recognized by anyone skilled in the art that other water-soluble polymers and other drugs may be utilized in a similar form of drug delivery. Examples of water-soluble polymers (denoted hereon by P) that can be used as carriers in such a drug delivery system are polyethylene glycols (PEG), polyvinyl alcohol, polyhydroxyethyl methacrylate, polyacrylamide, polyacrylic acid, polyethyloxazoline, polyvinyl pyrrolidinone, and polysaccharides such as chitosan, alginates, hyaluronic acid, dextrans, etc. These polymers can be covalently linked to the drugs by means of linkages (denoted hereon by X) such as ester, diester, urethane, amide, secondary or tertiary amine, ether etc.

The purpose of covalently linking a water-insoluble or poorly water-soluble drug (denoted hereon by D) to a water-soluble polymer is to solubilize the drug in water to enable its delivery in a soluble form into the body. The solubility of taxol in water is very low, approximately 0.03 mg/ml, and at the required dosage of 200-500 mg, this requires the infusion of a liter of fluid using a cremaphore to solubilize the drug. Thus it is desired to improve the solubility of taxol by conjugating it with a water-soluble polymer.

With reference to Figure 1, Table I below shows the chemical formulas of derivatives of taxol with linear PEGs and some intermediates used in the preparation of these derivatives. The R_1 and R_2 substituents on taxol vary

according to derivative and are indicated by the compound numbers which are used throughout the specification. Taxol itself is represented by compound 1.

Table I

5	Compound Number	R ₁ (2'-position)	R ₂ (7-position)
	1	OH	OH
	2	OCOC ₃ H ₃ N ₂	OCOC ₃ H ₃ N ₂
	3	OCONH(CH ₂ CH ₂ O) _n CH ₃	OCONH(CH ₂ CH ₂) _n CH ₃
10	4	OCOOCH ₂ CCl ₃	OH
	5	OCOOCH ₂ CCl ₃	OCONH(CH ₂ CH ₂ O) _n CH ₃
	6	OH	OCONH(CH ₂ CH ₂ O) _n CH ₃
	7	OOC(CH ₂) ₂ COO(CH ₂ CH ₂ O) _n CH ₃	OH
	8	OOC(CH ₂) ₂ COOH	OH
15	9	OOC(CH ₂) ₃ COOH	OH
	10	OOC(CH ₂) ₂ COO(CH ₂ CH ₂ O) _n CH ₃	OH
	11	OOC(CH ₂) ₃ COO(CH ₂ CH ₂ O) _n CH ₃	OH

n is the degree of polymerization or number of repeat units in the polymer chain and is dependant on the molecular weight of the polymer.

20 Table II below shows the general chemical formulas of linear PEG and its derivatives and intermediates utilized to obtain the drug-PEG conjugates. The general formula for PEG and derivatives is R₃(CH₂CH₂O)_nCH₂CH₂R₄ in which the flanking groups R₃ and R₄ vary according to the derivative. PEG itself is represented by compound 12 in which R₃ = R₄ = OH. Thus a 'linear bifunctional' PEG is represented by the compound 12 having two hydroxyl groups available for chemical reaction while a 'linear monofunctional' PEG is represented by compound 13 in which one end of the PEG is 'capped' with a nonreactive alkoxy or aryloxy group.

Table II

	Compound	Abbreviation	R ₃	R ₄
	12	PEG	HO	OH
5	13	MPEG	CH ₃ O	OH
	14	MPEG-amine	CH ₃ O	NH ₂
	15	MPEG-COOH	CH ₃ O	OCO(CH ₂) ₂ COOH
	16	PEG-monoacrylate	HO	OCOCH=CH ₂
	17	PEG-monomethacrylate	HO	OCOC(CH ₃)=CH ₂
10	18	MPEG-acrylate	CH ₃ O	OCOCH=CH ₂
	19	MPEG-methacrylate	CH ₃ O	OCOC(CH ₃)=CH ₂

The derivative of taxol having PEG at the 2' and 7 positions (compound 3) was prepared by first reacting taxol with 1,1-carbonyldiimidazole (CDI) as a coupling agent and subsequently with monomethoxy polyethylene glycol-amine (MPEG-amine, 14) to obtain the PEG derivative coupled by a urethane linkage which is relatively stable to hydrolysis. This derivative was not expected to have a high biological activity since the 2' position on taxol was substituted. In order to prepare a 7-PEG derivative of taxol, the 2' position was first protected with the [(2,2,2-trichloroethyl)oxy]carbonyl, or 'troc' protective group by reaction of taxol with 2,2,2-trichloroethyl chloroformate. The 2' protected derivative was then reacted with CDI and MPEG-amine as above to obtain the 2'-troc-7-PEG derivative followed by removal of the troc group to obtain the 7-PEG taxol. The water-solubility of these derivatives was determined in a UV spectrophotometer which clearly showed an absorbance for taxol when coupled to PEG. A wide range of MPEG-amine molecular weights, as well as other linear PEGs used for coupling of drugs can be utilized. Typically a molecular weight range of 200-100000 (corresponding n values between 5-2500) could be utilized for the derivatization. A preferred range is 600-20000 (n = 10-500) and the most preferred range is 1000-10000 (n =


20-250).

Another approach used was to deliver taxol in a soluble form such as the PEG derivative that could be hydrolyzed to release taxol in an active form after delivery of the drug-polymer conjugate. In this method taxol was linked to PEG at the 2' position by a readily hydrolyzable ester linkage. Two approaches were adopted to synthesize this derivative. The first involved modifying the hydroxyl end groups of MPEG with succinic anhydride to obtain the succinyl derivative of MPEG (15). This derivative was esterified with the 2'-hydroxyl on taxol using dicyclohexyl carbodiimide (DCC) and 4-dimethylamino pyridine (DMAP) to obtain the derivative 7. The second approach involved modification of the 2' hydroxyl on taxol with succinic or glutaric anhydride to obtain the succinyl (8) or glutaryl (9) derivative of taxol which was esterified with the MPEG hydroxyl using DCC and DMAP as before. Both these procedures resulted in the formation of 2'-MPEG taxol (10 or 11) that was readily hydrolyzable in an aqueous environment to give back active taxol and water-soluble carrier. A monofunctional PEG (MPEG) or a bifunctional PEG (regular PEG) could be used for this reaction. PEG (MPEG and/or PEG) molecular weights 200-100000 ($n = 5-2500$) could be utilized for the derivatization. A preferred range is 600-20000 ($n = 10-500$) and the most preferred range is 1000-10000 ($n = 20-250$).

In the above functionalization techniques, the number of drug molecules per carrier molecule is restricted to a maximum of two taxol molecules per molecule of PEG, and only one taxol per MPEG. In order to increase the number of taxols per carrier molecule, PEGs with multiple arms such as branched molecules or star molecules are used. A branched PEG was produced by solution polymerization of the monoacrylate derivative (16) or monomethacrylate

derivative (17) of PEG-2000 in the presence of the thermal free radical initiator, 2,2'-azobisisobutyronitrile (AIBN). Thus the number of available sites for coupling the drug to the 'brush-like' polymer was dependant on the number of PEGs having a free hydroxyl group that were incorporated into the growing polymer chain during the polymerization process. Any one of the reactions mentioned above for linear PEGs could be utilized to covalently link molecules of taxol to the branched polymer. Also, mixtures of PEG-monoacrylates of differing molecular weights could be utilized for the synthesis of a brush-like polymer in which the 'bristles' are of differing lengths. The general formula of branched PEGs synthesized for subsequent drug attachment is shown below:

15



20 wherein A = $\text{R}_5 (\text{CH}_2\text{CH}_2\text{O})_n \text{CH}_2\text{CH}_2\text{OCO}$, and $\text{R}_5 = \text{HO}$ (in case of PEG) or CH_3O (in case of MPEG);

wherein B = $\text{R}_5 (\text{CH}_2\text{CH}_2\text{O})_p \text{CH}_2\text{CH}_2\text{OCO}$, and $\text{R}_5 = \text{HO}$ (in case of PEG);

wherein $\text{R}_6 = \text{H}$ or CH_3 ;

25 wherein m, n, and p are the degrees of polymerization or number of repeat units in the polymer chain and are dependant on the molecular weight of the polymer. Typically m, n, and p = 5-2500.


When a drug D is bound to the above polymer through a covalent linking group X, the general formula remains the same except for R_5 which is replaced by D-X-.


'Star' molecules of PEG available commercially, e.g., 8-arm PEG, MW 22800, were functionalized with taxol using the techniques described above. These molecules may have a 'central core' of divinyl benzene (DVB) which is

anionically polymerized under controlled conditions to form living poly DVB nuclei having a predetermined number of active sites to which ethylene oxide is added to produce a known number of arms of PEG and quenched with water when the desired molecular weight is achieved. Alternately, they may have an oligomeric glycerol central core that is ethoxylated and used to initiate polymerization of ethylene oxide and quenched with water when the desired molecular weight is achieved. The range of usable molecular weights of these polymers ranges between 5000 and 200000 with a preferred range of 10000-100000 and a most preferred range of 20000-70000. The upper limit of this latter range is necessary for the carrier to be excreted from the circulatory system. A general formula for the star PEG is

15 $[\text{HO}-(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2]_q-(\text{central core})$ in which q is the number of arms of PEG attached to the central core. When a drug D is bound to the star polymer through a covalent linking group X , the general formula remains the same except for $\text{HO}-$ which is replaced by $D-X-$. The number of arms, q can

20 vary between 2 and 100. A schematic of a star PEG molecule with 8 arms ($q = 8$) is shown below:



wherein  = arms of PEG

and  = central core

A modified version of the branched polymer system also used as a drug carrier was synthesized by copolymerization of a PEG or MPEG monoacrylate with

'functional monomers'. Functional monomers are defined as those monomers that bear reactive functional groups that can be utilized for coupling of drugs. Examples of these are acrylic acid to provide carboxylic acid groups, allyl amine to provide primary amine groups, allyl alcohol to provide additional hydroxyl groups, or allyl chloride to provide chloride groups attached to the backbone of the branched copolymer. The copolymerization of acrylic acid and MPEG-5000 monoacrylate was carried out in toluene in the presence of the thermal free radical initiator, 2,2'-azobisisobutyronitrile (AIBN). Nonlimiting examples of functional monomers used in the synthesis are those bearing the carboxyl group, e.g., acrylic acid, vinyl acetic acid (3-butenic acid) and higher homologues; those bearing the amine group, e.g., allyl amine and higher homologues; those bearing the hydroxyl group, e.g., allyl alcohol (2-propene-1-ol) and higher homologues; allyl chloride (3-chloropropene), other unsaturated halides and corresponding higher homologues. The presence of these pendant functional groups allows for the attachment of a wide range of drugs possessing different functionalities. The general formula of copolymers of PEG and functional monomers synthesized for subsequent drug attachment is shown below:



wherein $\text{A} = \text{R}_5(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{OCO}$ and $\text{R}_5 = \text{HO}$ (for PEG) or CH_3O (for MPEG);

wherein $\text{R}_6 = \text{H}$ or CH_3 ;

wherein $\text{R}_7, \text{R}_8 = \text{H}, \text{CH}_3, \text{alkyl}, \text{or aryl}$;

wherein $\text{E} = \text{optionally alkyl or aryl}$;

wherein n and 4 are degrees of polymerization and vary between 5 and 2500 ;

and J = COOH, OH, CHO, NH₂, Cl, Br, or I.

When a drug D is bound to the above polymer at the site J, through a covalent linking group W (selected from the same groups as X, but not necessarily identical),
5 J is replaced by D-W-. Also the drug may be bound at A, in which case R₅ is replaced by D-X-.

The above text describes the production of taxol derivatives with water-soluble polymers. These soluble polymeric carriers containing the bound drug may be
10 crosslinked to produce an insoluble polymer matrix which is water-swellaable and has hydrogel properties. Such a matrix may be prepared in the form of a sphere, disc, cylinder, etc. that could be subsequently implanted at a suitable site for sustained release of the bound drug by hydrolysis.
15 Such a matrix is prepared by utilizing a branched or star PEG in which a portion of the available sites are functionalized by polymerizable acrylate or methacrylate groups and the remainder are bound to the drug. This polymer is isolated, dissolved in aqueous buffer (or
20 organic solvent) and crosslinked by a free radical process that may be thermally initiated or photoinitiated. Following crosslinking, the gel is desiccated by drying in vacuum and stored dry until before use when it is hydrated. To carry out the crosslinking step in organic solvent, the
25 polymer is dissolved at a suitable concentration to obtain a solution of mild viscosity, a thermal initiator such as AIBN, or a UV photoinitiator such as 2,2-dimethoxy-2-phenyl acetophenone (DMPA) is added. To prepare the gel in the form of a disk, the solution is poured into a mould and
30 heated or exposed to long wave UV radiation to crosslink the polymer. If the crosslinking step is to be carried out in aqueous medium, the same procedure is followed except for replacing the organic solvent with an aqueous buffer, adding a water-soluble UV initiator such as 2,2'-azobis-(2-
35 amidinopropane)hydrochloride (AAPH) and exposing to UV

light, or using a visible light initiated system comprising the dye ethyl eosin and cocatalyst triethanol amine and exposing the sample to green light in the region of 500-580nm. A small quantity of bifunctional crosslinkers may also be added, e.g., tetraethylene glycol diacrylate. The degree of substitution of available sites by polymerizable groups is varied depending on the degree of crosslinking and drug loading required. The presently preferred ratio of unsaturated groups to all available sites is between 0.04 and 0.75. A more preferred range is between 0.1 to 0.5.

The preferred mode of drug delivery for the soluble derivatives of taxol is intravenous. The polymer-drug conjugate is dissolved in normal saline or a physiological buffer and infused intravenously.

The invention will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLES

Example 1

Taxol-PEG: PEG at both 2' and 7 position coupled with a relatively stable urethane linkage.

10 mg (0.0117 mmol) of Taxol (Sigma Chemical) was dissolved in 5 ml of chloroform and a ten fold excess of 1,1-carbonyldiimidazole (CDI, 18.95 mg., 0.117 mmol) was added to activate the hydroxyl (OH) groups on the Taxol molecule. (There are three OH groups on Taxol (compound 1) at the 1, 7, and the 2' positions. The OH at the 2' position is the most labile followed by the OH at the 7 position while the OH at the 1 position is sterically hindered and nonreactive. This procedure resulted in the formation of the Taxol-CDI (2) derivative at the 2' and 7

positions). The reaction was allowed to proceed for 2 hours at room temperature and then extracted in water 3 times to remove the excess CDI and imidazole formed as a reaction product and then dried over anhydrous sodium sulfate. Monomethoxy polyethylene glycol amine (MPEG-amine, 14) of molecular weight 5000 g/mol was added (58.5 mg, 0.117 mmol) to the reaction mixture and allowed to react for 2 hours with stirring at room temperature. The product was extracted with 10% aqueous potassium nitrate, dried over anhydrous sodium sulfate and evaporated in vacuo or freeze dried.

UV spectroscopy (Shimadzu Instruments) was utilized to determine if Taxol had been derivatized and if a water-soluble PEG-Taxol derivative (3) substituted at the 2' and/or 7 position was obtained. Taxol was added to water (in an attempt to see any solubility that may result) to give a suspension which was filtered and a scan of this sample was obtained. No characteristic absorbances for taxol were seen implying a negligible solubility below the detection limit of the instrument. A scan of taxol dissolved in ethanol, however, showed the characteristic absorbance in the region of 240 nm and a shoulder around 270-280 nm. Next, MPEG-amine was dissolved in water and showed negligible absorption in the region of interest. A scan of the freeze dried product (PEG-taxol derivative) was added to water (100% of this product did not dissolve in water) and filtered. It showed the characteristic absorption of taxol indicating that the drug was solubilized in water as a result of the coupling reaction with PEG.

Example 2

Taxol-PEG: PEG at 7 position coupled with a relatively stable urethane linkage.

It is known that the C13 ester side chain and the 2'-hydroxyl group on the side chain are essential for biological activity (Mathew et al., 1992, *J. Med. Chem.* 35:145-151). The introduction of a substituent at the 2' position has resulted in a loss in the ability to promote microtubule assembly. Derivatives of taxol having a substituent at the 7 position however, retain their ability to alter cell proliferation and microtubule polymerization. The following procedure describes a method to obtain water-soluble PEG derivatives of taxol having the PEG substituent at the 7 position and not the 2' position.

The 2'-hydroxyl on taxol was first protected using the [(2,2,2-trichloroethyl)oxy]carbonyl, or 'troc' protective group. Taxol (50 mg) in chloroform (5 ml) and pyridine (0.1 ml) was cooled to -20°C and treated with 2,2,2-trichloroethyl chloroformate (0.008 ml) for 45 minutes. Workup by standard methods yielded the 2'-troc derivative (4) together with small amounts of taxol and 2', 7-bis troc taxol. The product could be isolated by TLC with ethyl acetate-hexane (1:1) as solvent: yield 85%.

The product 4 was reacted as in example 1 to obtain the 2'-troc-7-MPEG taxol (5). The protective troc group was then removed by dissolving 5 in 2 ml methanol-acetic acid (9:1) and addition of zinc dust (40 mg). The mixture was stirred for 10 minutes at room temperature, filtered to remove excess zinc and the 7-PEG taxol (6) obtained by precipitation with diethyl ether and drying in vacuo.

Example 3

Water-soluble Prodrugs of Taxol: PEG at the 2' Position Bound by a Hydrolyzable Linkage (Reaction of Taxol 2'-OH with PEG-COOH)

A PEG derivative of Taxol at the 2'-OH position will make taxol soluble in aqueous medium. An ester at this position will be hydrolyzed to give back taxol in its active form. This strategy is utilized in the delivery of
5 taxol in a water-soluble form. This form is called the 'prodrug'.

Preparation of MPEG-1900-COOH: MPEG-1900 (9.5g, 5 mmol) was dissolved in toluene (50ml) and dried by distilling off most of the toluene; succinic anhydride
10 (20g, 20mmol) was added and the mixture was stirred for 5 hours on an oil bath at 150°C. The mixture was cooled, taken up in dichloromethane and precipitated in ether. The product (15) was reprecipitated twice using dichloromethane/ether; yield 8.5g (85%). Homologues of
15 succinic anhydride, e.g., glutaric anhydride, may be utilized for this reaction.

Preparation of MPEG-Taxol Ester (7): MPEG-COOH (0.5 mmol COOH) and taxol (0.55mmol) were added to dichloromethane (20ml), and dicyclohexyl carbodiimide (DCC,
20 Aldrich, 0.65mmol), and 4-dimethyl aminopyridine (DMAP, Aldrich, 0.125mmol) were added as coupling agents. The mixture stirred for 3-4 hours at room temperature, the precipitate of dicyclohexyl urea (DCU) filtered and the filtrate evaporated to dryness in vacuum. The residue was
25 extracted with acetone and the product 7 precipitated by ether; yeilds 70-85%.

Example 4

*Water-Soluble Prodrugs of Taxol: PEG at the 2' Position Bound by a Hydrolyzable Linkage (Reaction of Taxol
30 2'-COOH with PEG-OH)*

An alternate strategy to the one in example 3 is to make a carboxylic acid derivative of taxol and esterify

it with untreated PEG, i.e., PEG having available OH groups. The carboxylic acid derivatives of taxol are prior art and have been synthesized by esterification with succinic or glutaric anhydrides (Deutsch et al., 1989; *J. Med. Chem.* 32:788-792).

Preparation of 2'-Succinyl Taxol (8): Taxol (0.5g, 0.59 mmol) and succinic anhydride (0.90g, 7.6mmol) in 12 ml of pyridine were allowed to react for 3 hours at room temperature after which the mixture was evaporated to dryness *in vacuo*. The residue was treated with 20ml of water, stirred for 20min and then filtered. The precipitate was dissolved in acetone, water slowly added, and the fine crystals of 8 were collected; yield (75-85%).

Preparation of 2'-Glutaryl Taxol (9): A similar procedure used for 8 by reaction of taxol with glutaric anhydride gave 85-95% yield of the derivative 9 when recrystallized from chloroform/benzene.

Preparation of MPEG-Taxol Ester (10, 11): Essentially the identical procedure used for preparation of product 7 was used here, the MPEG-COOH being replaced by untreated MPEG (13) and taxol by products 8 or 9 to obtain respectively the products 10 and 11. Yields obtained were in the range of 75-85%. The product 10 was identical to the product 7 obtained in example 3.

Homologues of 8 and 9 obtained by reaction of taxol with homologues of cyclic anhydrides such as succinic anhydride may be utilized for preparation of PEG-Taxol esters.

Example 5

Water-Soluble Taxol-PEG Derivatives: Use of Branched Chain or 'Star' PEGs for a Multiplicity of

Attachment Sites for Taxol.

The use of derivatives in examples 1 through 4 describes the covalent attachment of taxol to PEGs that are monofunctional (1 available attachment site per molecule) such as MPEG or at most, those that are bifunctional (regular PEGs, 2 available attachment sites per molecule). In order to increase the efficiency of drug delivery in terms of increasing the ratio of the mass of drug to the mass of inactive carrier (in this case PEG), it is desirable to utilize a molecule (carrier) that for a given molecular weight, can carry several molecules of drug as opposed to one or two molecules per molecule of carrier.

Preparation of Branched PEG-Taxol: Branched polymers of PEG produced by polymerization of a monoacrylate or monomethacrylate derivative were also used for covalent attachment of taxol. The monoacrylate (16) and monomethacrylate (17) derivatives of PEG (Macrochem Labs) of molecular weights ranging from 1000 to 20000 having one free hydroxyl end group were polymerized in solution to obtain branched polymers of a given molecular weight. 10g (5mmol) of PEG 2000 monoacrylate was dissolved in 100 ml of dry toluene. A thermal free radical initiator 2,2'-azobis isobutyronitrile (AIBN) was added 0.016g (0.1 mmol) and the solution heated to 80°C. The reaction was allowed to proceed overnight and the polymer precipitated from solution with diethyl ether. The polymer was further purified by redissolution in toluene and reprecipitation with ether and dried *in vacuo*. The branched PEG so obtained was then coupled to taxol by the techniques illustrated in examples 1-4. Those skilled in the art will recognize that other free radical initiators or initiating mechanisms, and other PEG derivatives, e.g., vinyl ethers of PEG, may be utilized to obtain branched molecules.

Preparation of Star PEG-Taxol: 'Star' PEG specifically 8-arm PEG, MW 22800 (Macrochem Labs) was used as obtained. This PEG had 8 potential coupling sites per molecule. Other such PEGs may also be used with a greater
5 number of arms and higher molecular weight preferably less than 100000 to allow clearance from the body by the kidneys. The reactions described in examples 1 through 4 were utilized to covalently attach drugs such as taxol to these molecules.

10

Example 6

Water-Soluble Taxol-PEG Derivatives: Use of Copolymers of PEG and Functional Monomers for a Multiplicity of Attachment Sites for Taxol.

Another strategy for drug delivery is the
15 copolymerization of functional monomers with polymerizable derivatives of PEG such as MPEG-acrylate (18) or methacrylate (19) or the corresponding 'mono' derivatives of regular PEG. Examples of functional monomers used in the synthesis are those bearing the carboxyl group, e.g.,
20 acrylic acid, vinyl acetic acid (3-butenic acid) and higher homologues; those bearing the amine group, e.g., allyl amine and higher homologues; those bearing the hydroxyl group, e.g., allyl alcohol (2-propene-1-ol) and higher homologues; allyl chloride (3-chloropropene), other
25 unsaturated halides and corresponding higher homologues, and unsaturated compounds bearing aldehyde groups. The presence of these pendant functional groups allows for the attachment of a wide range of drugs possessing different functionalities.

30

Copolymerization of MPEG-acrylate and Acrylic Acid: MPEG-acrylate was prepared by reaction of MPEG (Nippon Oil and Fat) with acryloyl chloride (Aldrich). MPEG-5000 (10g, 2mmol) was dissolved in toluene (150 ml)

and approximately 50 ml of toluene was distilled over to ensure removal of water. Acryloyl chloride (4mmol) that had been distilled immediately prior to use, was added after the MPEG/toluene had been cooled on an ice bath.

5 Triethylamine (4mmol, not essential) was added and the reaction mixture refluxed for 4 hours. Triethylamine hydrochloride, formed as a by product was filtered and the MPEG-acrylate precipitated with by addition of excess ether to the filtrate. The product was purified by

10 reprecipitation from toluene and then dried *in vacuo*: Yield 90%.

Acrylic acid was vacuum distilled prior to use. MPEG-acrylate (1mmol) and acrylic acid (1mmol) were dissolved in dry toluene and the free radical thermal

15 initiator AIBN (0.05mmol) added. The reaction mixture was heated to 80°C, the reaction was allowed to proceed overnight and the polymer precipitated from solution with diethyl ether. The polymer was further purified by redissolution in toluene and reprecipitation with ether and

20 dried *in vacuo*. The copolymer containing labile carboxylic acid groups was then coupled to taxol by the technique described for product 7.

Example 7

Hydrogels Containing Bound Taxol for Sustained

25 *Release Drug-delivery.*

Star (8-arm) PEG-22800 (2.85g, containing 1 mmol OH groups) was dissolved in 25 ml of dry toluene. The solution was cooled on an ice bath and 0.33mmol of freshly distilled acryloyl chloride was added and the reaction

30 mixture kept at 70°C for 4 hours. The toluene was removed by vacuum distillation and the partial acrylate derivative redissolved in 25ml of dry dichloromethane. 2'-succinyl taxol (0.8 mmol, prepared as in example 4), DCC (0.95 mmol)

and DMAP (0.18 mmol) were added to the reaction mixture which was stirred at room temperature for 3-4 hours. The precipitate of DCU was filtered and the PEG-partial acrylate derivatized with taxol (star-PEG-acrylate-taxol, SPAT) was precipitated in excess ether and dried in vacuo. Yield: 70-85%

This derivative could be crosslinked by free radical polymerization in organic solvent or in aqueous medium by addition of the appropriate initiators. In organic solvent, the UV photoinitiator DMPA was used. A 10 wt% solution of SPAT in dichloromethane containing 0.05% DMPA was poured in an open stainless steel mold in the shape of a small disc 1cm diameter and thickness 0.5cm and exposed to long-wave UV radiation from a mercury arc lamp. The solution gelled within 30 seconds. The gel was freeze dried. When placed in water of aqueous medium, the gelled discs would swell and imbibe water.

A similar crosslinking procedure was utilized in aqueous buffer except the photoinitiator in this case was water-soluble UV initiator AAPH at 0.05% concentration. Other organic soluble initiators such as benzil, and other aqueous photoinitiating systems such as the ethyl eosin, triethanol amine system were also successful.

A branched PEG polymer, or a branched copolymer could be utilized instead of the star polymer to achieve a similar drug immobilized gel.

CLAIMS

What is claimed is:

1. A Taxol derivative having the general formula R_1-T-R_2 :
 - 5 wherein $R_1 = X-P$ or $-OH$, and R_1 is located on the 2'-carbon of the taxol side chain;
 - wherein $R_2 = -X-P$ or $-OH$, and R_2 is located on the 7-carbon atom of the ring structure;
 - but R_1 and R_2 are not simultaneously OH ;
 - 10 and the drug taxol T , is covalently linked to a water-soluble polymer P , through a covalent linking group X .
2. The composition of claim 1 in which said covalent linking group X is selected from ester, diester,
 - 15 urethane, amide, secondary or tertiary amine, or ether groups.
3. The composition of claim 1 in which said water-soluble polymer P is based on polyethylene glycol.
4. The composition of claim 3 in which said
 - 20 polyethylene glycol is a linear monofunctional polyethylene glycol, wherein:
 $R_1 = R_9(CH_2CH_2O)_nCH_2CH_2X-$ or $-OH$, and R_9 is selected from alkoxy or aryloxy;
 - $R_2 = -R_1$ or $-OH$;
 - 25 and n is a number between, and including, 5 and 2500.

5. The composition of claim 3 in which the polyethylene glycol is a linear bifunctional polyethylene glycol, wherein:

- 5 $R_1 = R_{10}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{X}-$, or $-\text{OH}$, and $R_{10} = \text{T-X}$ -
where X is at the 2'-position on the taxol side chain;
 $R_2 = -\text{OH}$ or $R_{11}(\text{CH}_2\text{HC}_2\text{O})_n\text{CH}_2\text{CH}_2\text{X}-$, and $R_{11} = \text{T-X}$ -
where X is at the 7-position on the taxol ring structure;
10 where R_1 and R_2 do not simultaneously comprise a polyethylene glycol;
and n is a number between, and including, 5 and 2500.

6. The composition of claim 3 in which the
15 polyethylene glycol is a branched polyethylene glycol, and the taxol-polyethylene glycol derivative has the general formula:



- wherein $\text{A} = \text{R}_{12}(\text{CH}_2\text{CH}_2\text{O})_n\text{CH}_2\text{CH}_2\text{-Y-}$ and $\text{R}_{12} = \text{HO-}$, or alkoxy, or aryloxy, or T-X- ;
25 wherein $\text{B} = \text{T-X-(CH}_2\text{CH}_2\text{O)}_p\text{CH}_2\text{CH}_2\text{-Y-}$;
wherein $\text{R}_6 = -\text{H}$ or $-\text{CH}_3$;
wherein X is located at either the 2' position of the taxol side chain or the 7 position of the taxol ring structure but not both simultaneously;
30 wherein Y is a covalent linking group independently selected from the same groups as X;
and m, n, and p are numbers between, and including, 5 and 2500.

7. The composition of claim 3 in which the polyethylene glycol is a star polyethylene glycol, and the taxol-polyethylene glycol derivative has the general formula;

5 [T-X-(CH₂CH₂O)_nCH₂CH₂]_q-(central core)
wherein X is located at either the 2' position of
the taxol side chain or the 7 position of the
taxol ring structure but not both simultaneously;
wherein q is the number of arms of a linear
10 polyethylene glycol attached to a central core;
wherein n is a number between, and including, 5
and 2500;
and q is a number between, and including, 2 and
100.

15 8. The composition of claim 3 in which said water-soluble polymer is a copolymer of a linear polyethylene glycol and a functional monomer, the resultant copolymer-taxol derivative having the general formula;



wherein A = $R_{12}(CH_2CH_2O)_nCH_2CH_2-Y-$ and $R_{12} = HO-$, or alkoxy, or aryloxy, or T-X-;

wherein $R_6 = -H$ or $-CH_3$;

wherein Y is a covalent linking group independently selected from the same groups as X;

wherein R_7 and R_8 are selected from H, or alkyl, or aryl;

wherein E is optionally selected from alkyl or aryl;

wherein $G = -W-T$, wherein W is a covalent linking group independently selected from the same groups

as X, and T is linked to E through W.
wherein W is located at either the 2' position of
the taxol side chain or the 7 position of the
taxol ring structure but not both simultaneously;
and r is a number between, and including, 5 and
2500.

5
9. The composition of claim 6 in which a
fraction of taxol substituents are replaced with
unsaturated groups capable of undergoing free radical
10 polymerization.

10. The composition of claim 7 in which a
fraction of taxol substituents are replaced with
unsaturated groups capable of undergoing free radical
polymerization.

15
11. The composition of claim 8 in which a
fraction of taxol substituents are replaced with
unsaturated groups capable of undergoing free radical
polymerization.

12. The composition of claim 6 in the form of a
20 crosslinked insoluble gel containing covalently bound
taxol.

13. The composition of claim 7 in the form of a
crosslinked insoluble gel containing covalently bound
taxol.

25
14. The composition of claim 8 in the form of a
crosslinked insoluble gel containing covalently bound
taxol.

15. A water-soluble drug-polymer conjugate having the general formula P-X-D;
wherein the polymer P is water-soluble;
wherein the drug D, is by itself water-insoluble or poorly water-soluble;
and the drug and polymer are linked by the covalent linkage X.


16. The composition of claim 15 in which said water-insoluble or poorly water-soluble drug D is taxol or amphotericin B.

17. The composition of claim 15 in which said water-soluble polymer is crosslinked to form an insoluble gel.

18. A method of solubilizing a water-insoluble or poorly water-soluble drug by attaching to a water-soluble polymeric carrier by a covalent linkage.

19. The method of 18 in which said water-insoluble or poorly water-soluble drug is taxol or amphotericin B.

20. A water-soluble drug delivery system comprising water-soluble polymers of polyethylene glycol conjugated to a drug D by a covalent linking group, the drug-polymer conjugates having the general formulas:



wherein the water-soluble polymer is a branched polyethylene glycol;
wherein said drug is linked to said water-soluble

29

polymer by said covalent linking group denoted by X;

wherein $A = R_5(CH_2CH_2O)_nCH_2CH_2-Y-$ and $R_5 = HO-$, or alkoxy, or aryloxy, or D-X-;

5 wherein $B = D-X-(CH_2CH_2O)_pCH_2CH_2-Y-$;

wherein $R_6 = -H$ or $-CH_3$;

wherein Y is independently selected from the same groups as X;

10 wherein m, n, and p are numbers between, and including, 5 and 2500;

OR



20 wherein the water-soluble polymer is a branched copolymer of a linear polyethylene glycol and a functional monomer;

wherein said drug is linked to said water-soluble polymer by covalent linking groups X or W;

25 wherein A and R_6 are defined as above;

wherein R_7 and R_8 are selected from $-H$, $-CH_3$, alkyl, or aryl;


wherein E is optionally selected from alkyl or aryl;

30 and $G = -W-D$, wherein W is a covalent linking group independently selected from the same groups as X, and D is linked to E through W;

wherein r is a number between, and including 5 and 2500;

OR

35



wherein the water soluble polymer is a star polyethylene glycol;

wherein said drug is linked to said water-soluble polymer by said covalent linking group denoted by X;

wherein q is the number of arms of a linear polyethylene glycol attached to a central core and q is a number between, and including, 2 and 100;

and n is defined as above.


21. The composition of claim 20 wherein said drug D is water-insoluble or poorly water-soluble.

22. The composition of claim 21 as a means of solubilizing said water-insoluble or poorly water-soluble drug in water.

23. The composition of claim 21 in which said water-insoluble or poorly water-soluble drug is taxol or amphotericin B.

1/1

FIGURE 1



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/05344

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : CO7D 305/14,
US CL : 549/510, 549/511

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : NONE

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS, taxol, Polyethylene glycol

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,534,899 (Sears), 13 Aug. 1985	1-23
Y	US, A, 5,059,699 (Kingston et al.), 22 Oct., 1991.	1-23
Y	"Polymer Preprints", 31, 1990, Nathan et al., Polyethylene Glycol-Lysine Copolymer: New Biocompatible Polymers For Biomedical Application", pp. 213-214.	1-23
Y	"Eur. Polym. J.", Vol. 19, No. 12, 1983, Zalipsky et al., Attachment of Drugs to Polyethylene Glycols, pp. 1177-1183	1-23

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A		document defining the general state of the art which is not considered to be part of particular relevance
* E		earlier document published on or after the international filing date
* L		document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
* O		document referring to an oral disclosure, use, exhibition or other means
* P		document published prior to the international filing date but later than the priority date claimed
	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
	* &	document member of the same patent family

Date of the actual completion of the international search

09 AUGUST 1993

Date of mailing of the international search report

09 SEP 1993

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

WARREN IVY

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US93/05344

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	"J. Med. Chem.", 24, 1981, Cecchi et al., Synthesis and Pharmacological Evaluation of Poly (oxyethylene) Derivatives of 4-Isobutyl phenyl-2-propionic Acid (Ibuprofen), pp. 622-625	1-23
Y	"J. Med. Chem.", Vol. 16, No. 5, 1973. Zilkha et al., Polyethylene Glycol Derivatives of Procaine, pp. 573-574.	